

EM Basic- Female Specific Abdominal Pain

(This document doesn't reflect the views or opinions of the Department of Defense, the US Army or the SAUSHEC EM residency, © 2011 EM Basic, Steve Carroll DO. May freely distribute with proper attribution)

History

First question- is the patient pregnant?

- Urine HCG and UA often done from triage
- If delay in getting urine, get an IV and check a serum qualitative HCG

OPQRST questions about pain

Onset, Provocation, Quality, Radiation, Severity, and Time

Associated signs and symptoms

Nausea/vomiting/diarrhea, back pain, urinary symptoms

Female specific questions

Missed periods, vaginal bleeding, discharge

Gs and Ps

G= Pregnancies P= Live births- know them for OB consultation

Medical history

Special attention to surgical history, history of ectopic pregnancies or spontaneous/elective abortions

Sexual history

- Is the patient sexually active currently or in the past?
- (Kick parents out of room for younger patient)
- Monogamous relationship?
- History or concern for STDs?

Exam

Don't dive for the abdomen- do an HEENT exam, heart/lung exam

Uncover the abdomen and ask patient to point where it hurts the most

- Top of iliac crest- more often abdominal cause of pain
- Lower towards inguinal ligament- pelvic pathology
- Not absolute but can help you guide your workup

Differential Diagnosis

- Ectopic Pregnancy
- Threatened abortion/miscarriage
- Normal pregnancy
- STDs- Gonorrhea, Chlamydia, Trichomonas
- Pelvic Inflammatory Disease
- Tubo-Ovarian abscess

Ovarian Torsion

Ovarian Cyst

Bacterial Vaginosis

Vaginal Candidiasis (BV and Candidiasis don't cause pain but are frequently encountered in the workup)

Workup

UA and HCG (looking for UTI, pregnancy status)

CBC (check for WBC- low yield and anemia)

Chem 10 (standard part of abdominal pain workup)

LFTs

Lipase

Coags (these may not be necessary in every patient- let your clinical judgement be your guide but appy often overlaps with pelvic pathology)

Pelvic Exam

GC/Chlamydia

Wet Prep/KOH

Quantitative HCG (if the patient is pregnant)

Pelvic Exam tips

Do an external exam to look for lesions or obvious discharge

As you insert the speculum, examine for lesions or discharge

GC and Chlamydia swab- insert in cervix and twist 360 degrees

Wet Prep/KOH- swab vaginal walls

Bimanual exam

Touching the cervix is uncomfortable- do this slowly

Say to the patient "I know this is uncomfortable but does this hurt when I move my fingers" and look for reaction.

If the patient seems comfortable this is NOT cervical motion tenderness

If the patient is in a lot of distress, then this is CMT

Check the adnexa

Palpate the abdomen while checking adnexa to see where the patient has the most pain

Imaging

PEARL- HUGE overlap between pelvic and abdominal pathology- keep your differential open

Transvaginal Pelvic ultrasound- helpful in suspected ectopic pregnancy (covered in a later podcast). Should see IUP at HCG of approximately 1,500

Ovarian Torsion- “Classic”- Sudden onset of sharp stabbing pain in lower abdomen with nausea or vomiting.

PEARL- A negative pelvic ultrasound DOES NOT rule out torsion

Intermittent torsion is possible

Time sensitive diagnosis- ovary salvage rate greatly declines after 4 hours

High suspicion= OB/GYN consult and admission for observation
Most torsions have large cysts but not all

PEARL- If radiologist doesn't want to do TVUS after hours because “an ultrasound doesn't rule out torsion”, say you are concerned for tubo-ovarian abscess or large hemorrhagic cyst

Ovarian Cyst- Cyst on same side as pain with good blood flow to ovary
Can be discharged with pain control, OB/GYN followup

STDs

Treatment depends on how fast your GC/Chlamydia test is

Send-out = Empirically treat all patients with discharge and/or CMT

PEARL- ALWAYS treat for gonorrhea and chlamydia

(Often co-exist and missing it can have long term fertility consequences)

Cervicitis treatment (no CMT but positive swab)

Ceftriaxone 125mg IM

1 gram of azithromycin PO x1 dose

(PCN or cephalosporin allergic- 2 grams azithromycin PO x1)

PID treatment (CMT and/or discharge)

Ceftriaxone 250mg IM

Doxycycline 100mg PO BID for 14 days

PEARL- Admit patients who are ill appearing, pregnant, unresponsive to outpatient treatment, PO intolerant

Caution patient to not have sex for seven days after treatment stopped and to get partner tested to avoid re-infection

Fitz-Hugh Curtis- PID infection that has spread to the liver capsule and causes perihepatitis- may have RUQ pain and tenderness, right shoulder pain. LFTs may be elevated but not always. Treatment is to treat for PID

Tubo-Ovarian Abscess- non-specific symptoms- abdominal/pelvic pain, history of STDs, severe or recurrent PID. Usually untreated PID that has formed an abscess. Diagnosed with TVUS, treated with IV abx, surgical drainage

Trichomonas- Motile organisms on Wet Prep. Sxs- itching, discharge, dysuria dyspareunia

Treatment- Flagyl 2 grams PO x1 or 500mg PO BID for 7 days

Treat partner as well- men are often asymptomatic

No alcohol while on flagyl (disulfaram reaction = vomiting)

Bacterial Vaginosis- not an STD but frequently diagnosed- overgrowth of Gardnerella Vaginalis. Sxs- malodorous discharge. Clue cells on Wet Prep

Treat all symptomatic patients AND pregnant patients

Treatment- Flagyl 2 grams PO or 500mg PO BID for 7 days

Candidiasis- “yeast infection” usually after abx. Fungal elements on Wet Prep.

Treatment- Fluconazole- 150mg PO x1 or topical treatments

Topical treatments can control sxs better, can use with fluconazole

PEARL- with all pelvic swabs- false negatives occur so treat clinically for what you think is the most serious cause

How to approach positive STD tests- Sometimes women will be suspicious for STDs, sometime not. Have a frank discussion of what you found and what it means. Talk to the patient alone- if you have to, devise a ruse to get anyone else out of the room (go get the car, go to the pharmacy, sign paperwork, etc.) Make sure the patient knows that they have an STD but let them know that they could have been asymptomatic for a long time and that you aren't pointing fingers. Let them know the importance of completing the full course of treatment and following up with their OB/GYN or primary care doctor. Let them know that you aren't accusing anyone of cheating. Make sure to stress to the patient the importance of getting their partner tested and treated as well. If you are treating PID empirically (send-out GC/Chlamydia) make sure that the patient knows this and they understand that you are treating them for an STD without knowing whether or not they have one. Make sure you have a good phone number to call the patient back with a positive test result.

Contact- steve@embasic.org